

Catherin Donaldson ACPIN 2009  
**Reference list**

- Bobath B. Adult hemiplegia: evaluation and treatment. 3rd ed. London: Heinemann Publishing 1990
- Bohannon RW. Muscle strength and muscle training after stroke: review article. *J Rehabil Med* 2007;39:14-20
- Bohannon RW, Warren ME, Cogman KA. Motor Variables Correlated with the Hand-to-mouth Maneuver in Stroke Patients. *Arch Phys Med Rehabil* 1991;72:682-684
- Bonita R, Beaglehole R. Recovery of motor function after stroke. *Stroke* 1988;19:1497-1500
- Butefisch C, Hummelsheim H, Denzler P, Mauritz K-H. Repetitive training of isolated movements improves the outcome of motor rehabilitation of the centrally paretic hand. *Journal of the Neurological Sciences* 1995; 130:59-68
- Broeks JG, Lankhorst GJ, Rumping K, Prevo AJH. The long term outcome of arm function after stroke: results of a follow-up study. *Disability and Rehabilitation* 1999;21:357-364
- Carr J, Shepherd R. Stroke Rehabilitation: Guidelines for Exercise and Training to Optimize Motor Skill. Butterworth Heinemann 2003
- Dean CM, Shepherd RB. Task-related training improves performance of seated reaching tasks after stroke; A randomised controlled trial. *Stroke* 1997; 28(4):722-728
- Donaldson C, Tallis R, Miller S, Sunderland A, Lemon R, Pomeroy V. Effects of Conventional Physical Therapy and Functional Strength Training on Upper Limb Motor Recovery after Stroke: a Randomised Phase II Study. *Neurorehabil Neural Repair*. 2008 Dec 24.
- Donaldson C, Tallis R, Pomeroy V. A treatment schedule of conventional physical therapy provided to enhance upper limb sensori-motor recovery after stroke: expert criterion validity and intra-rater reliability. *Physiotherapy* 2009 January 27
- Duncan PW, Goldstein LB, Horner RD, Landsman PB, Samsa GP, Matchar DB. Similar Motor Recovery of Upper and Lower Extremities after Stroke. *Stroke* 1994;25:1181-1188
- Duncan P, Studenski S, Richards L, Gollub S, Min Lai S, Reker D, Perera S, Yates J, Koch V, Rigler S, Johnson D. Randomised clinical trial of therapeutic exercise in subacute stroke. *Stroke* 2003; 34:2173-2180
- Elliott J M and Connolly K J. A Classification of Manipulative Hand Movements. *Developmental Medicine and Child Neurology* 1984 26: 283-296.
- Feys H, De Weerd W, Verbeke G, Steck G, Capiou C, Kiekens C, Dejaeger E, Van Hoydonck G, Vermeersch G, Cras P. Early and repetitive stimulation of the arm can substantially improve the long-term outcome after stroke: a 5-year follow-up study of a randomised trial. *Stroke* 2004; 35(4):924-929
- Gowland C, deBruin H, Basmajian JV, Plews N, Burcea I. Agonist and antagonist activity during voluntary upper-limb movement in patients with stroke. *Phys Ther* 1992; 72(9):624-629
- Hachisuka K, Umezue Y, Ogata H. Disuse muscle atrophy of lower limbs in hemiplegic patients. *Arch Phys Med Rehabil* 1997;78(1):13-18
- Hendricks H T, van Limbeek J, Geurts A C, Zwarts M J. Motor recovery after stroke: a systematic review of the literature. *Arch Phys Med Rehabil* 2002;83:1629-1637
- Hendricks HT, Zwarts MJ, Plat EF, van Limbeek J. Systematic review for early prediction of motor and functional outcome after stroke by using motor-evoked potentials. *Arch Phys Med Rehabil* 2002;83:1303-1308

- Ho Jang S, Kim Y-H, Cho J-H, Park J-W, Kwon Y-H. Cortical reorganization induced by task-oriented training in chronic hemiplegic stroke patients. *NeuroReport* 2003;14(1):137-141
- Hochstenbach J, Mulder T. Neuropsychology and the relearning of motor skills following stroke. *International Journal of Rehabilitation Research*. 1999;22:11-19
- Hunter SM, Crome P, Sim J, Donaldson C, Pomeroy VM. Development of Treatment Schedules for research: a systematic review to identify methodologies used and worked example of 'mobilisation and tactile stimulation' for stroke patients. *Physiotherapy* 2006;92:195-207
- Johansen-Berg H, Dawes H, Guy C, Smith SM, Wade DT, Matthews PM. Correlation between motor improvements and altered fMRI activity after rehabilitative therapy. *Brain* 2002;125:2731-2742
- Jones M, Tallis R, Pomeroy V. Physiotherapists may not practice what is implied from treatment "labels". *Age and Ageing*. 2003;32(2):229-30
- Kwakkel G, Van Peppen R, Wagenaar RC, Dauphinee SW, Richards C, Ashburn A, Miller K, Lincoln N, Partridge C, Wellwood I, Langhorne P. Effects of augmented exercise therapy time after stroke. A meta-analysis. *Stroke* 2004;35:2529-2536
- Kwakkel G, Kollen BJ, van der Grond J, Prevo AJ. Probability of regaining dexterity in the flaccid upper limb: impact of severity of paresis and time since onset in acute stroke. *Stroke*. 2003;34(9):2181-6
- Levy CE, Nichols DS, Schmalbrock PM, Keller P, Chakeres DW. Functional MRI Evidence of Cortical Reorganization in Upper-Limb Stroke Hemiplegia Treated with Constraint-Induced Movement Therapy. *Am J Phys. Med. Rehabil* 2001;80(1):4-12
- Liepert J, Graef S, Uhde I, Leidner O, Weiller C. Training-induced changes of motor cortex representations in stroke patients. *Acta Neurol Scand* 2000b;101:321-326
- Liepert J, Bauder H, Miltner WHR, Taub E, Weiller C. Treatment-Induced Cortical Reorganisation after Stroke in Humans. *Stroke* 2000a;31:1210-1216
- Liepert J, Bauder H, Miltner WHR, Bauder H, Sommer C, Dettmers, Taub E, Weiller C. Motor cortex plasticity during constraint-induced movement therapy in stroke patients. *Neuroscience letters* 1998; 250:5-8
- Norris CM. Sports injuries: diagnosis and management. Second Edition. Butterworth-Heinemann, Oxford. 1998.
- Parker VM, Wade DT, Langton-Hewer R. Loss of arm function after stroke: measurement, frequency and recovery. *Int Rehabil Med* 1986;8:69-73
- Parry RH, Lincoln NB, Vass CD. Effect of severity of arm impairment on response to additional physiotherapy early after stroke. *Clinical Rehabilitation* 1999;13:187-198
- Patten C, Lexell J, Brown HE. Weakness and strength training in persons with poststroke hemiplegia: Rationale, method and efficacy. *Journal of Rehabilitation Research and Development* 2004;41(3):293-312
- Pomeroy VM, Tallis RC. Physical Therapy to improve movement performance and functional ability poststroke. Part 1. Existing evidence. *Reviews in Clinical Gerontology* 2000 A;10:261-290
- Pomeroy VM, Tallis RC. Need to focus research in stroke rehabilitation. *The Lancet* 2000 B;355:836-837
- Robertson IH, Ridgeway V, Greenfield E, Parr A. Motor recovery after stroke depends on intact sustained attention: a 2-year follow-up study. *Neuropsychology*. 1997;11:290-295

- Taixeira-Salmela LF, Olney SJ, Nadeau S, Brouwer B. Muscle strengthening and physical conditioning to reduce impairment and disability in chronic stroke survivors. *Arch Phys Med Rehabil* 1999;80:1211-1218
- Taub E, Miller NE, Novack TA, Cook III EW, Fleming WC, Nepomuceno CS. Technique to improve chronic motor deficit after stroke. *Arch Phys Med Rehabil* 1993;74:347-54
- Thielman GT, Dean CM, Gentile AM. Rehabilitation of reaching after stroke: task-related training versus progressive resistive exercise. *Arch Phys Med Rehabil* 2004;85:1613-1618
- Van der Lee JH, De Groot V, Beckerman H, Wagenaar RC, Lankhorst GJ, Bouter LM. The intra- and interrater reliability of the Action Research Arm Test: A practical test of upper extremity function in patients with stroke. *Arch Phys Med Rehabil* 2001B;82:14-19
- Van der Lee H, Snels IAK, Beckerman H, Lankhorst GJ, Wagenaar RC, Bouter LM. Exercise therapy for arm function in stroke patients: a systematic review of randomised controlled trials. *Clin Rehabil* 2001;15:20-31
- Van Peppen RP, Kwakkel G, Wood-Dauphinee S, Hendriks HJ, Van der Wees PJ, Dekker J. The impact of physical therapy on functional outcomes after stroke: what's the evidence? *Clinical Rehabilitation*. 2004;18(8):833-62
- Winstein CJ, Rose DK, Tan SM, Lewthwaite R, Chui HC, Azen SP. A randomised controlled comparison of upper extremity rehabilitation strategies in acute stroke: A pilot study of immediate and long-term outcome. *Arch Phys Med Rehabil* 2004; 85:620-628
- Yoo EY, Chung BI. The effect of visual feedback plus mental practice on symmetrical weight-bearing training in people with hemiparesis. *Clin Rehab* 2006;20(5):388-97